



CD40LG gene

CD40 ligand

Normal Function

The *CD40LG* gene provides instructions for making a protein called CD40 ligand, which is found on the surface of immune system cells known as T cells. CD40 ligand attaches like a key in a lock to its receptor protein, CD40, which is located on the surface of immune system cells known as B cells. B cells are involved in the production of proteins called antibodies or immunoglobulins that help protect the body against infection. There are several classes of antibodies, and each one has a different function in the immune system. B cells are able to mature into the cells that produce immunoglobulin M (IgM) without any signals from other cells. In order for B cells to mature into the cells that produce antibodies of a different class, the CD40 receptor must interact with CD40 ligand. When these two proteins are connected, they trigger a series of chemical signals that instruct the B cell to start making immunoglobulin G (IgG), immunoglobulin A (IgA), and immunoglobulin E (IgE).

CD40 ligand is also necessary for T cells to interact with other cells of the immune system, and it plays a key role in T cell differentiation (the process by which cells mature to carry out specific functions).

Health Conditions Related to Genetic Changes


X-linked hyper IgM syndrome

More than 150 mutations in the *CD40LG* gene have been found to cause X-linked hyper IgM syndrome. These mutations lead to the production of an abnormal CD40 ligand or prevent production of this protein. If CD40 ligand does not attach to its receptor on B cells, these cells cannot produce IgG, IgA, or IgE antibodies. Mutations in the *CD40LG* gene also impair the T cell's ability to differentiate and interact with immune system cells. People with X-linked hyper IgM syndrome are more susceptible to infections because they do not have a properly functioning immune system.

Chromosomal Location

Cytogenetic Location: Xq26.3, which is the long (q) arm of the X chromosome at position 26.3

Molecular Location: base pairs 136,648,177 to 136,660,390 on the X chromosome (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- CD40 antigen ligand
- CD40L
- CD40L_HUMAN
- CD154
- gp39
- hCD40L
- HIGM1
- IGM
- IMD3
- T-B cell-activating molecule
- T-BAM
- TNF-related activation protein
- TNFSF5
- TRAP
- tumor necrosis factor (ligand) superfamily member 5

Additional Information & Resources

Educational Resources

- Immunobiology (fifth edition, 2001): Isotype switching requires expression of CD40L by the helper T cell and is directed by cytokines
<https://www.ncbi.nlm.nih.gov/books/NBK27142/#A1192>

GeneReviews

- X-Linked Hyper IgM Syndrome
<https://www.ncbi.nlm.nih.gov/books/NBK1402>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28CD40LG%5BTIAB%5D%29+OR+%28CD40+ligand%5BTIAB%5D%29%29+AND+%28%28tumor+necrosis+factor+ligand+superfamily+member+5%5BMAJR%5D%29+OR+%28cd40+ligand%5BMAJR%5D%29+OR+%28cd154%5BMAJR%5D%29+OR+%28tnf+superfamily,+member+5%5BMAJR%5D%29+OR+%28cd40l%5BMAJR%5D%29%29+AND+%28Genes%5BMH%5D%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+1800+days%22%5Bdp%5D>

OMIM

- CD40 LIGAND
<http://omim.org/entry/300386>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
http://atlasgeneticsoncology.org/Genes/GC_CD40LG.html
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=CD40LG%5Bgene%5D>
- HGNC Gene Family: CD molecules
<http://www.genenames.org/cgi-bin/genefamilies/set/471>
- HGNC Gene Family: Endogenous ligands
<http://www.genenames.org/cgi-bin/genefamilies/set/542>
- HGNC Gene Family: Tumor necrosis factor superfamily
<http://www.genenames.org/cgi-bin/genefamilies/set/781>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=11935

- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/959>
- UniProt
<http://www.uniprot.org/uniprot/P29965>

Sources for This Summary

- Bhushan A, Covey LR. CD40:CD40L interactions in X-linked and non-X-linked hyper-IgM syndromes. *Immunol Res.* 2001;24(3):311-24. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/11817328>
- Durandy A. Hyper-IgM syndromes: a model for studying the regulation of class switch recombination and somatic hypermutation generation. *Biochem Soc Trans.* 2002 Aug;30(4):815-8. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/12196205>
- GeneReview: X-Linked Hyper IgM Syndrome
<https://www.ncbi.nlm.nih.gov/books/NBK1402>
- Gilmour KC, Walshe D, Heath S, Monaghan G, Loughlin S, Lester T, Norbury G, Cale CM. Immunological and genetic analysis of 65 patients with a clinical suspicion of X linked hyper-IgM. *Mol Pathol.* 2003 Oct;56(5):256-62.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/14514918>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1187335/>
- Jain A, Atkinson TP, Lipsky PE, Slater JE, Nelson DL, Strober W. Defects of T-cell effector function and post-thymic maturation in X-linked hyper-IgM syndrome. *J Clin Invest.* 1999 Apr;103(8):1151-8.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/10207167>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC408278/>
- Notarangelo LD, Lanzi G, Peron S, Durandy A. Defects of class-switch recombination. *J Allergy Clin Immunol.* 2006 Apr;117(4):855-64. Review.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16630945>
- Thusberg J, Vihinen M. The structural basis of hyper IgM deficiency - CD40L mutations. *Protein Eng Des Sel.* 2007 Mar;20(3):133-41. Epub 2007 Feb 16.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17307885>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/CD40LG>

Reviewed: October 2008
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services